



Immunization Schedule

	Birth	1 Month	2 Months	4 Months	6 Months	9 Months	12 Months	15 Months	18 Months	2 Years	4 Years	9-10 Years	11-12 Years	13-15 Years	16 Years and Older
Hepatitis B	#1	#2 ^a				#3									
Diphtheria, Tetanus, and Pertussis (DTaP)			#1	#2	#3			#4			#5				
Haemophilus influenzae type B (HiB)			#1	#2	#3			#4							
Inactivated Poliovirus (IPV)			#1	#2	#3			#4			#5				
Pneumococcal (PCV)			#1	#2	#3		#4								
Rotavirus (oral)			#1	#2	#3										
Influenza					#1										
Measles, Mumps, and Rubella (MMR)							#1 ^c				#2 ^d				
Varicella							#1 ^c				#2 ^d				
Hepatitis A									#1	#2					
Meningococcal													#1		#2
Tetanus, Diphtheria, and Pertussis (TDaP)													#1		
HPV (Gardasil)													v ^e	v ^e	v ^e
Meningococcal B															v ^f

a - Hepatitis B #2 must be given at least 28 days after Hepatitis B #1.

b - Must be at least 6 months of age to receive the influenza vaccine. If under 9 years of age and receiving first dose every, patient will need a booster dose within the same season after 30 days.

c - Must be on or after 1st birthday.

d - Must be on or after 4th birthday.

e - HPV vaccine is recommended for girls and boys 9 years of age and older. If patient begins HPV series before the age of 15, only two doses are needed at least 6 months apart. If administered after age 15, patient will need 3 doses with the first and second being 2 months apart and the second and third being 4 months apart.

f - Meningococcal B vaccine is recommended for girls and boys within this age. The series is administered in 2 doses at least 6 months apart.

The above schedule also corresponds to our recommended schedule for physicals.

Physicals are also recommended at 30 months of age. Annual physicals are recommended yearly for patients 3 years of age and older.